OmegaRx Zone, 2002, Kapitel "When the Brain goes wrong", Unterkapitel "Depression", S.120, 3 Seiten:

"Clinical depression is a disabling condition

[Dieses Kapitel habe ich NICHT übersetzt und AUCH NICHT in elektronischer Form zur Verfügung] [am Ende dieses Word-Dokuments habe ich aber die LITERATUR für das o.a. Kapitel angehängt,] [dort nach "depression" suchen

Anti-Inflammation Zone, 2005, Kapitel "Brain Drain Due To Silent Inflammation", Unterkapitel "Depression", S.279, 2 Seiten

2005-03-11

"Post-partum diet"

Dear Dr. Sears,

I am having a hard time losing my pregnancy weight post partum and am nursing a three-month-old. Are nursing mothers able to diet using the Zone and still ensure the baby will get everything he needs?

Because you are nursing, you will need some extra calories, so just add one block of each macronutrient to every meal and snack. Also make sure you are supplementing your diet with ultra-refined EPA/DHA concentrates (about 2.5 grams per day) to maintain adequate levels in the breast milk **and to prevent any post- partum depression**.



Fish oil may help relieve stubborn depression

Last Updated: 2002-10-17 10:00:24 -0400 (Reuters Health) NEW YORK (Reuters Health) -Daily supplements of an omega-3 fatty acid--found in fish and fish oil--may help alleviate the symptoms of depression in patients who do not respond to standard antidepressant medications, new research findings suggest. Dr. Malcolm Peet of the Swallownest Court Hospital in Sheffield, England and his colleague found that depressed patients who received a daily dose of 1 gram of an omega-3 fatty acid for 12 weeks experienced a decrease in their symptoms, such as sadness, anxiety and sleeping problems. The only side effect of the treatment appeared to be gastrointestinal problems, which Peet and his co-author Dr. David F. Horrobin of Laxdale Research, Ltd. in Stirling, Scotland, deemed "mild." All of the patients had tried other medications before enrolling in the current study, including selective serotonin reuptake inhibitors (SSRIs) such as Prozac and medications from an older family of drugs called tricyclic antidepressants. Both types of drug are considered standard treatments for depression. This is not the first study to suggest that omega-3 fatty acids, such as the form of eicosapenaenoic acid (EPA) used in this report, may help patients with psychiatric disorders. Previous researchers have suggested that the balance of omega-3 fatty acids in the brain may become skewed in people with depression, and earlier studies have shown that fish oil supplements can help alleviate the symptoms of schizophrenia and bipolar disorder, or manic depression. In addition, researchers have found that people who are depressed, as well as those diagnosed with cardiovascular diseases and other conditions associated with depression, have relatively low levels of omega-3 fatty acids in their blood. In the current study, reported in the October issue of the Archives of General Psychiatry, Peet and Horrobin asked 70 depressed patients who had not benefited from previous treatments to take a daily dose of either 1 gram, 2 grams or 4 grams of EPA, or an inactive drug. The treatment lasted 12 weeks. The investigators found that people given the 1 gram daily EPA dose experienced improvements--relative to those given the inactive drug--in all of the measured aspects of depression, including sadness, anxiety, low libido and suicidal tendencies. In fact, 69% of the patients treated with the 1-gram daily dose achieved a 50% reduction in their symptoms of depression, a result seen in only 25% of the patients given an inactive drug. "The effect of ethyl-eicosapentaenoate (the form of EPA used) applies to all major components of the depressive syndrome and is seen equally in the patient and physician assessments," the authors write. Peet and Horrobin did not note any improvements in the patients given higher doses of

the fatty acid relative to the placebo group, which they suggested may be due to the small number of people who were given either 2 grams or 4 grams per day. "Although there appeared to be a trend toward significant efficacy at the 4-gram per day dosage, larger studies would be required to elucidate possible beneficial effects of the higher dosages," they write.

Dr. Sears' Comments:

One reason they may have been having trouble getting a dose-response curve is that the fish oil preparation they used did not contain any DHA.

Fish oil helps ease depression

It works! Just like the book. After a serious case of depression (I've been diagnosed as bipolar), and at the encouragement of a doctor, I took on the Zone way of eating -- again. Food as an anti-depressant? That was a new one for me. I had tried the Zone once before, but because of a hectic work life with a lot of travel, I could never stay committed. This time was different. I stayed diligent. Within three weeks, I began to notice something different: Gone was my constant craving for fried food. (I'm a sucker for fried anything.) I started losing weight, or so I thought as I don't have a scale. Turned out it was even better. I was losing inches. Had a doctor's appointment recently and discovered I had only lost 10 pounds. That's surprising because I've dropped two inches in my waist and am wearing 34-inch waist jeans again ... and those are starting to look a little big. I feel so light. And I feel stronger. But wait, it gets better.

In November I saw the good Doc's new book, "The Omega Rx Zone." It was a fascinating read, but I must admit to being a bit sceptical. But he was right about the food stuff, so I ordered some fish oil, and all I can say is wow! I do think better. I do have better control of my emotions. I'm writing again (hooray!). I've been going for hour walks for some time now. But since taking the fish oil, I find myself walking much faster and for much longer. Every member of my family, including a dozen or so friends, were wiped out just prior to the holidays by a nasty chest cold. I got it, too, I guess. Sure felt like it, but within a day, it was dissipating, and two days later gone. Everyone else suffered with this cold for two weeks. My poor wife had to go on antibiotics. One of our close friends developed pneumonia from this cold. I felt a little guilty I felt so good, but also grateful. I love my fish oil. I found that approximately 10 grams a day is perfect for me. While much higher than a "normal" daily dose, 10 grams has a noticeable impact on my depression. I'm a customer for life! And given my new-found health, it sure seems like I'll be a customer for a long, long, long time. Thanks, Doc. I can't thank you enough.

Jim W.

Today's question

"Fish oil for depression"

Dear Dr. Sears, An earlier response addressed supplementing the zone diet with high-dose fish oil to treat depression. What is the exact dose recommended? Davidcwhite

Dear Davidcwhite: I usually recommend a minimum of four to six grams of fish oil per day. High dose fish oil would be between 10 and 20 grams per day. That would provide between 3 and 6 grams of long-chain Omega-3 fatty acids. Studies published by Harvard Medical School on the fish oil and depression used the equivalent of 30 grams of fish oil per day which would provide about 10 grams of long-chain Omega-3 fatty acids.



Dr. Barry Sears

Today's question

"Depression/eating disorders"

"Depression/eating disorders" Dear Dr. Sears, I want to know about the omega supplement and how it might help depression and eating disorders? Nancy

Dear Nancy, Fish oils rich in omega-3 fatty acids have been very beneficial to help treat depression. The dosage used in studies at Harvard Medical School was 10 grams of long-chain omega-3 fatty acids per day. At these doses, I would only recommend pharmaceutical-grade fish oils that have exceptional purity with regard to mercury, PCB, and other toxic contaminants.

Bang Som

Dr. Barry Sears

Today's question

"Fish oil & depression"

Dear Dr. Sears, I know this question has been asked many times, but I just wanted to get an idea of the exact amount and exact type of fish oil to take daily for the treatment of moderate depression and over-all health. I just bought a bottle of fish oil concentrate 1000mg. The bottle says 2 capsules contain 600mg Omega-3 Polyunsaturates, which is 360mg EPA and 240mg DHA. I've started taking 12 capsules daily. I feel slightly better already. Also, is there any way to get the pharmaceutical strength fish oil until you will sell it through your web site? Thank you. Sincerely, Shawn

Dear Shawn: The data from Harvard Medical School indicated that 9.6 grams of long-chain Omega-3 fatty acids per day had a significant effect on depression. The amount of Omega-3 fatty acids that you are currently taking is 7.2 grams. The type of fats that you are using is one-step below pharmaceutical grade. One company that makes pharmaceutical-grade fish oil is Omega Natural Health. Their web site is www.omegabrite.com.

Bay Son

Dr. Barry Sears

Dear Dr. Sears,
I am an alcoholic, as well as a long-time sufferer of severe depression , anxiety disorder, panic disorder, and OCD. (My daughter, 8, shows symptoms of these as well, including severe ADHD, which hasn't been helped by medication.) I take 999 mg of Campral three times a day as prescribed for the treatment of alcoholism, as well as 50 mg. of Paxil. Recently, after reviewing the data from the "OmegaRx Zone," I've redoubled my efforts to eat in the Zone and have upped my fish oil supplementation from six pills a day to one tablespoon a day. My question: Is the fish oil in any way going to adversely affect my brain while on Campral? There's no data for this that I can find regarding chemical interaction. Also, is giving my daughter one tablespoon of refined fish oil a day going to hurt her at only 60 pounds?
ALB
The fish oil will provide significant benefits to aid in the treatment of your existing neurological conditions. Furthermore, the fish oil will reduce many of the gastric side effects of Campral. Our studies with children used two tablespoons per day

2005-07-26

"Campral and fish oil"

Anxiety"

Does pharmaceutical-grade fish oil help anxiety?

of fish oil, so the dosage for your daughter is, if anything, a little on the low side.

Wendell

Dear Wendell,

BR> Virtually all neurological disorders respond very positively to high-dose fish oil supplementation. Although there are no clinical studies on treating anxiety specifically, the conditions that co-exist with anxiety, such as depression, are known to significantly benefit from high-dose fish oil.

Bay Son

Chapter 10--When the Brain Goes Wrong

Adams P, Lawson S, Sanigorski A, and Sinclair AJ. "Arachidonic acid to eicosapentaenoic acid ratio in blood correlates positively with clinical symptoms of depression." Lipids 31: S157-S161 (1996)

Agren JJ, Tormala ML, Nenonem MJ, and Haainea OO. "Fatty acid composition of erythrocyte, platelet, and serum lipids in strict vegetarians." Lipids 30: 365-369 (1995)

Agostoni C, Trojan S, Bellu R, Riva E, Bruzzese MG, and Giovannini M. "Development quotient at 24 months and fatty acid composition of the diet in early infancy." Arch Dis Child 76: 421-424 (1997)

Ahmann PA, Waltonen SJ, Olson KA, Theye FW, van Erem AJ, and LePlant RJ." Placebo-controlled evaluation of Ritalin side effects." Pediatrics 91: 1101-1106 (1993)

Aisen PS. "Anti-inflammatory therapy for Alzheimer's disease." Neurobiol Aging 21: 447-448 (2000)

Akiyama H, Arai T, Kondo H, Tanno E, Haga C, and Ikeda K. "Cell mediators of inflammation in the Alzheimer disease brain." Alzheimer Disease and Associated Disorders 14: S47-S53 (2000)

Aisen PS. "Anti-inflammatory therapy for Alzheimer's disease: implication of the prednisone trial." Acta Neurol Scand 176: 85-89 (2000)

Alvarez JC, Cremniter D, Lesieur P, Gregoire A, Gilton A, Macquin-Mavier I, Jarreu C, and Spreux-Varoquaux O. "Low blood cholesterol and low platelet serotonin levels in violent suicide attempters." Bio Psychiatry 45: 1066-1069 (1999)

Amen DG. Change Your Brain, Change Your Life. Random House. New York, NY (1998)

Amen DG. Healing ADD. GP Putnam. New York, NY (2001)

Attvall S, Fowelin J, Lager I, Von Schenck H, and Smith U. "Smoking induces insulin resistance--a potential link with the insulin resistance syndrome."

J Intern Med 233: 327-332 (1993)

Blaylock RL. Excitotoxins. Health Press, Santa Fe, NM (1995)

Bourre JM. "Function of polyunsaturated fatty acids in the nervous system." Prostglandins Leukot Essent Fatty Acids 48: 5-15 (1993)

Burdge GC, Wright SM, Warner JO, and Postle AD. "Fetal brain and liver phospholipid fatty acid composition in a guinea pig model of fetal alcohol syndrome: effect of maternal supplementation with tuna oil." J Nutr Biochem 8: 438-444 (1997)

Stevens LJ, Zentall SS, Deck JL, Abate ML, Watkins BA, Lipp SR, and Burgess JR "Essential fatty acid metabolism in boys with attention deficit-hyperactivity disorder." Am J Clin Nutr 62: 761-768 (1995)

Burgress JR, Stevens L, and Peck L. "Long-chain polyunsaturated fatty acids in children with attention-deficit hyperactivity disorder." Am J Clin Nutr 71: 327S-330S (2000)

Bush G, Frazier JA, Rauch SL, Seidman LJ, Whalen PJ, Jenike MA, Rosen BR, and Biederman J. "Anterior cingulated cortex dysfuction in attentiondeficit/hyperactivity disorder revealed by fMRI and the counting stroop." Biol Psychiarty 45: 1542-1552 (1999)

Carlson S and Werkman A. "A randomized trial of visul attention of preterm infants fed docosahexaenoic acid until two months." Lipids 31: 85-90 (1996)

Carrie I, Clement M, De Javel D, Frances H, and Bourre JM. "Learning deficits in the first generation OF1 mice deficient in (n-3) polyunsaturated fatty acids do not result from visual alteration." Neurosci Lett 266: 69-72 (1999)

Centonze D, Calabresi P, Giacomini P, and Berardi G. "Neurophysiology of Parkinson's disease: from basic research to clinical correlates." Clin Neurophysiology 110: 2006-2013 (1999)

Connor WE, Neuringer M, and Lin DS. "Dietary effects on brain fatty acid composition: the reversibility of n-3 fatty acids deficiency and turnover of docosahexaenoic acid in the brain erthrocytes, and plasma of rhesus monkeys." J Lipid Res 31: 237-247 (1990)

Connor WE, Neuringer M, and Reisbick S. "Essential fatty acids: importance of n-3 fatty acids in the retina and brain." Nutr Rev 50: 21-29 (1992)

Cooper NR, Kalaria RN, McGeer PL, and Rogers J. "Key issues in Alzheimer's disease inflammation." Neurobiology Aging 21: 451-453 (2000)

Conquer JA, Tierney MC, Zecevic J, Bettger WJ, and Fisher RH. "Fatty acid analysis of blood plasma of patients with Alzheimer's disease, other types of dementia, and cognitive impairment." Lipids 35: 1305-1312 (2000)

Delion S, Chalon S, Guilloteau D, Besnard JC, and Durand G. "Alpha-linolenic acid deficiency alters age-related changes of dopaminergic and serotoninergic neurotransmitters in the rat frontal cortex." J Neurochem 66: 1582-1591 (1996)

Delion S, chalon S, Herault J, Guilloteau D, Besnard JC, and Durand G. "Chronic dietary alpha-linolenic acid deficiency alters dopaminergic and serotoninergic neurotransmitters in rats." J Nutr 124: 2466-2476 (1994)

Ensel M, Milon H, and Malnoe A. "Effect of low intake of n-3 fatty acids during development of brain phospholipid, fatty acid composition and exploratory behavior in rats." Lipids 26: 203-208 (1991)

FentonWS, Hibbeln J, and Knable M. "Essential fatty acids, lipid membrane abnormalities, and the diagnosis and treatment of schizophrenia." Bio Psychiarty 47: 8-21 (2000)

Fernstrom JD. "Effects of dietary polyunsaturated fatty acids on neuronal function." Lipids 34: 161-169(1999)

Freychet P. "Insulin receptors and insulin actions in the nervous system." Diabetes Metab Res Rev 16: 390-392 (2000)

Gayo A, Mozo L, Suarez A, Tunon A, Lahoz C, and Gutierrez C. "Inteferon beta treatment modulates TNF and interferon gamma spontaneous gene expression in MS." Neurology 52: 1764-1770 (1999)

Gallai V, Sarchielli P, Trequattrini A, Franceschini M, Floridi A, Firenze C, Alberti A, Di Benedetto D, and Stragliotto E. "Cytokine secretion and eicosanoid production in the peripheral blood mononuclear cells of MS patients undergoing dietary supplementation with n-3 polyunsaturated fatty acids." J Neuroimmunol 56: 143-153 (1995)

Gibson RA, Neuman MA, and Makrides M. "Effect of dietary docosahexenoic acid on brain compostion and neural function in term infants." J Lipid Res 34: S177-S181 (1996)

Glueck CJ, Tieger M, Kunkel R, Tracy T, Speirs J, Streicher P, and Illig E. "Improvement in symptoms of depression and in an index of life stressors accompany treatment of severe hypertriglyceridemia." Biol Psycharity 34: 240-252 (1993)

Hamazaki T, Sawazaki S, Itomura M, Asaoka E, Nagao Y, Nishimura N, Yazawa K, Kuwamori T, and Kobayashi M. "The effect of docosahexaenoic acid on aggression in young adults." J Clin Invest 97: 1129-1134 (1996)

Hamaszki T, Sawazaki S, Itomura M, Nagao Y, Thienprasert A., Nagasawa T, and Watanabe S. "Effect of docosahexaenoic acid on hostility." World Rev Nutr Diet 88: 47-52 (2001)

Harvey BH and Bouwer CD. "Neuropharmacology of paradoxic weight gain with selective serotonin reuptake inhibitors." Clinica Neuropharm 23: 90-97 (2000)

Hibbeln JR and Salem N. "Dietary polyunsaturated fatty acids and depression: when cholesterol does not satisfy." Am J Clin Nutr 62: 1-9 (1995)

Hibbeln JR. "Fish consumption and major depression." Lancet 351: 1213 (1998)

Hibbeln JR. "Seafood consumption and homicide mortality." World Rev Nutr Diet 88: 41-46 (2001)

Hohlfeld R and Wiendl H. "The ups and downs of multiple sclerosis therapeutics." Ann Neurology 49: 281-284 (2001)

Holman RT, Johnson SB, and Ogburn PL. "Deficiency of essential fatty acids and membrane fluidity during pregnancy and lactation." Proc Natl Acad Sci USA 88: 4835-4839 (1991)

Horrobin DF. "Essential fatty acids, prostaglandins, and alcoholism: an overview." Alcohol Clin Exp Res 11: 2-9 (1987)

Hoozemans JJM, Rozemuller AJM, Janssen I, De Groot CJA, Veerhuls R, and Eikelenboon P. "Cyclooxygenase expression in microglia and neurons in Alzheimer's disease and control brain." Acta Neuropathol 101: 2-8 (2001)

Hoozemans JJM, Veerhuis R, Janssen I, Rozemuler AJM, and Eikelenboom P. "Interleukin-1 beta induced cyclooxygenase 2 expression and prostaglandinE2 secretion by human neroblastoma cells: implications for Alzheimer's disease." Exp Gerontology 36: 559-570 (2001)

Holden RJ, Pakula IS, and Mooney PA. "The role of brain insulin n the neurophysiology of serious mental disorders: review." Med Hypothesis 52: 193-200 (1999)

Ichiyama T, Okada K, Lipton JM, Matsubara T, Hayashi T, and Furukawa S. "Sodium valproate inhibits production of TNF and IL-6 and activation of NF-kappa B." Brain Res 857: 246-251 (2000)

Ikemoto A, Nitta A, Furukawa S, Ohishi M, Nakamure A, Fujii Y, and Okuyama H. "Dietary n-3 fatty acid deficiency decreases nerve growth factor content in rat hippocampus." Neurosci Lett 285: 99-102 (2000)

Iso H, Stampfer MJ, Manson JE, Rexrode K, Hu F, Hennekens CH, Colditz GA, Speizer FE, and Willett WC. "Prospective study of fat and protein intake and risk of intraparenchymal hemorrhage in women." Circulation 103: 856-863 (2001)

Kalmijn S, Foley D, White L, Burchfiel CM, Curb JD, Petrovitch H,Ross GW, Havlik RJ, and Launer LJ. "Metabolic cardiovascular syndrome and risk of dementia in Japanese-American elderly men." Arterioscler Thromb Vasc Biol 20: 2255-2260 (2000)

Kademi M, Wallstrom E, Andersson M, Piehl F, Di Marco R, and Olsson T. "Reduction of both pro- and anti-inflammatory cytokines after 6 months of interferon beta-1a treatment of multiple sclerosis." J Neurochem 103: 202-210 (2000)

Kawas CH and Brookmeyer R. "Aging and the public health: effects of dementia." N Engl J Med 344 1160-1161 (2001)

Knapp HR, Reilly IAG, Alessandrini P, and FitzGerald GA. "In vivo indexes of platelet and vascular function during fish-oil administration in patients with atherosclerosis." N. Engl J Med 314: 937-942 (1986)

Kyle DJ, Schaefer E, Patton G, and Beiser A. "Low serum docosahexaenoic acid is a significant risk factor for Alzheimer's dementia." Lipids 34: S245 (1999)

Lauritzen I, Blondeau N, Heurteaux C, Widmann C, Romey G, and Lazdunski M. "Polyunsaturated fatty acids are potent neuroprotectors." EMBO J 19: 1784-1793 (2000)

Lauritzen L, Hansen HS, Jorgensen MH, and Michaelsen KF. "The essentially of long chain n-3 fatty acids in relation to development and function of the brain and retina." Prog Lipid Res 40: 1-94 (2001)

Maes M. "Fatty acid composition in major depression: decreased n-3 fractions in cholesterol esters and increased C20:n6/C20:5n3 ratio in cholesterol ester and phospholipids." J Affect Dis 38: 35-46 (1996)

Maes M, Christophe A, Delanghe J, Altamura C, Neels, H, and Meltzer HY. "Lowered omega-3 polyunsaturated fatty acids in serum phospholpids and cholesteryl esters of depressed patients." Pscyhiarty Res 85: 275-291 (1999)

Manev H, Tolga U, Sugaya K, and Qu T. "Putative role of neuronal 5-lipoxygenase in an aging brain." FASEB J 14: 1464-1469 (2000)

Mayeux R, Costa R, Bell K, Merchant C, Tung MX, and Jacobs D. "Reduced risk of Alzheimer's disease among individuals with low caloric intake." Neurology 59: S296-S297 (1999)

McGeer PL, Shulzer M, and McGeer EG. "Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: a review of 17 epidemiological studies." Neurology 47: 425-432 (1996)

McGeer PL, McGeer EG, and Yasojima K. "Alzheimer disease and neuroinflammation." J Neural Transm 59: 53-57 (2000)

Mills DE, Prkochin KM, Harvey KA, and Ward RP. "Dietary fatty acid supplementation alters stress reactivity and performance in man." J Human Hypertension 3: 111-116 (1989)

Minami M, Kimura S, Endo T, Hamaue N, Hirafuji M, Togashi H, Matsujoto M, Yoshika M, Saito H, Watanabe S, Kobayashi T, and Okuyama H. "Dietary docosahexaenoic acid increases cerebral acetylcholine levels and improves passive avoidance performance in stroke-prone spontaneously hypertensive rats." Pharmacol Biochem Behav 58: 1123-1129 (1997)

Miyanga K, Yonemura K, Takagi T, Kifune R, Kishi Y, Miyakawa F, Yazawa K, and Shirota Y. "Clinical effects of DHA in demented patients." J Clin Ther Med 11: 881-901 (1995)

Mischoulon D and Fava M. "Docosahexanoic acid and omega-3 fatty acids in depression." Psychiatr Clin North Am 23: 785-794 (2000)

Montine TJ, Sidell KR, Crews BC, Markesbery WR, Marnett LJ, RobertsLJ, and Morrow JD. "Elevated CSF prostaglandin E2 levels in patients with probable AD." Neurology 53: 1495-1498 (1999)

Moriguchi T, Greiner RS, and Salem N Jr. "Behavioral deficits associated with dietary induction of decreased brain docosahexaenoic acid concentration." J Neurochem 75: 2563-2573 (2000)

Moses H and Sriram S. "Interferon beta and the cytokine trial. Where are we going?" Neurology 52: 1729-1730 (1999)

Nagatsu T, Mogi M, Ichinose H, and Togari A. "Cytokines in Parkinson's disease." J Neural Transm 58: 143-151 (2000)

Nelson GJ, Schmidt PS, Bartolini GL, Kelley DS, and Kyle D. "The effect of dietary docosahexaenoic acid on platelet function, platelet fatty acid composition, and blood coagulation in humans." Lipids 32: 1129-1136 (1997)

Neuroinflammation Working Group. "Inflammation and Alzheimer's disease." Neurobiology Aging 21: 383-421 (2000)

Nightingale S, Woo E, Smith AD, French JM, Gale MM, Sinclair HM, Bates D, and Shaw DA "Red blood cell and adipose tissue fatty acids in active and inactive multiple sclerosis patients." Acta Neurol Scand 82: 43-50 (1990)

Nordvik I, Myhr K-M, Nyland H, and Bjerve KS. "Effect of dietary advice and n-3 supplementation in newly diagnosed MS patients." Act Neurol Scand 102: 143-149 (2000)

Pasinetti GM and Aisen PS. "Cyclooxygenase -2 expressin is increased in frontal cortex of Alzheimer's disease brain." Neuroscience 87: 319 324 (1997)

Pawlosky RJ and Salem N. "Ethanol exposure causes a decrease in docosahexanenoic acid and an increase in docosapentaenoic acid in feline brain and retina." Am J Clin Nutr 61: 1284-1289 (1995)

Peet M. "Essential fatty acid deficiency in erthrocyte membranes from chronic schizophrenic patients and clinical effects of dietary supplementation." Pros Leukotrienes Essential Fatty Acids 55: 71-75 (1996)

Peet M, Brind J, Ramchand CN, Shah S, and Vankar GK.

"Two double-blind placebo-controlled pilot studies of eicosapentaenoic acid in the treatment of schizophrenia." Schizophr Res 49: 243-251 (2001)

Pratico D and Rojanowski JQ. "Inflammatory hypothesis: novel mechanisms of Alzheimer's neurodegradation and new therapeutic targets?" Neurobiology Aging 21: 441-445 (2000)

Rasmuson S, Andrew R, Nasman B, Seckl JR, Walker BR, and Olsson T. "Increased glucocorticoid production and altered cortisol metabolism in women with mild to moderate Alzheimer's disease." Bio Psychiarty 49: 547-552 (2001)

Remarque EJ, Bollen ELEM, Weverling-Rijnsburger AWE, Laterveer JC, Blauw GJ, and Westendorp RGJ. "Patients with Alzheimer's disease display a pro-inflammatory phenotype." Exp Gerontology 36: 171-176 (2001)

Reisbick S, Neuringer M, Hasnain R, and Connor WE. "Home cage behavior of rhesus moneky with long-term deficiency of omega-3 fatty acids." Physiol Behav 55: 231-239 (1994)

Riviere S, Biroluez-Aragon I, and Vellas B. "Plasma protein glycation in Alzhemier's disease." Glycocongjugate J 15: 1039-1042 (1998)

Sachdev P. "Attention deficit hyperactivity disorder in adults." Psychological Med 29: 507-514 (1999)

Shoulson I. "DATATOP: a decade of neuroprotective inquiry. Parkinson Study Group. Deprenyl and tocopherol antioxidative therapy of Parkinsonism." Ann Neurol 44: S160-166 (1998)

Sinclair AJ and Crawford MA. "The effect of a low fat maternal diet on neonatal rats." Br J Nutr 29: 127-137 (1973)

Sonderberg M, Edlund C. Kristensson K, and Dallner G. "Fatty acid composition of brain phospholipids in aging and Alzheimer's disease." Lipids 26: 421-423 (1991)

Stein J. "The neurobiology of reading difficulties." Prostaglandins Leuko Essen Fatty Acids 63: 109-116 (2000)

Stevens LJ and Burgess J. "Omega-3 fatty acids in boys with behavior, learning, and health problems." Physiology Behavior 59: 915-920 (1996)

Stevens LJ, Zentall SS, Deck JL, Abate ML, Watkins BA, Lipp SA, and Burgess JR. "Essential fatty acid metabolism in boys with attention-deficit hyperactivity disorder." Am J Clin Nutr 62: 761-768 (1995)

Stewart WF, Kawas C, Corrada M, and Metter EJ. "Risk of Alzhemier's disease and duration of NSAID use." Neurology 48: 626-632 (1997)

Stoll AL. The Omega-3 Connection. Simon and Shuster. New York, NY (2001)

Stoll AL and Severus E. "Mood stabilizers: shared mechanisms of action at post-synaptic signal transduction and kindling process." Harvard Rev of Psychiarty 59: 915-920 (1999)

Stoll AL, Sverus E, freeman MP, Rueter S, Zhoyan HA, Diamond E, Cress KK, and Marangell LB. "Omega-3 fatty acids in bipolar depression: a preliminary double-blind, placebo-controlled trial." Arch Gen Psychiarty 56: 407-412 (1999)

Seung Kim HF, Weeber EJ, Sweatt JD, Stoll AL, and Marangell LB. "Inhibitory effects of omega-3 fatty acids on protein kinase C activity in vitro." Mol Psychiatry 6: 246-248 (2001)

Stordy BJ. "Benefit of docosahexaenoic acid supplements to dark adaption in dyslexics." Lancet 346: 385 (1995)

Tanner CM. "Dopamine agonists in early therapy. Promise and problems." JAMA 284: 19711973 (2000)

Tanskanen A. "Fish consumption, depression, and suicidality in a general population." Arch Gen Psychiarty 58: 512-513 (2001)

Tatton WG, Wadia JS, Ju WY, Chalmers-Redman RM, and Tatton NA. "Deprenyl reduces neuroal apoptosis and faciliates neuronal outgrowth by altering protein synthesis without inhibiting monoamine oxidase." J Neural Transm 48: 45-59 (1996)

Taylor KE and Richardson AJ. "Visual function, fatty acids and dyslexia." Prostglandins Leuko Essen Fatty Acids 63: 89-93 (2000)

Terano T, Fujishiro S, Ban T, Ymamoto K, Tanaka T, Noguchi Y, Tamura Y, K Yazawa, and Hirayama T. "Docosahexaenoic acid supplementation improves moderately severe dementia from thromobotic cerebrovascular diseases. "Lipids 34: S345-S346 (1999)

Uauy R, Peirano P, Hoffman D, Mena P, Birch D, and Birch E. "Role of essential fatty acids in the function of the developing nervous system." Lipids 31: S167-S176 (1996)

Venters HD, Dantzer R, and Kelly KW. "A new concept in neurodegeneration: TNF is a silencer of survival signals." Trends in Neuroscience 23: 175-180 (2000)

Virkkunen ME, Horrobin DF, Douglas K, Jenkins K, and Manku MS. "Plasma phospholipid essential fatty acids and prostaglandin in alcholic, habitually violent, and impulsive offenders." Bio Psych 22: 1087-1096 (1987)

Vitkovic L, Bockaert J, and Jacque C. "Inflammatory cytokines: neuromodulators in normal brain?" J Neurochem 74: 457-471 (2000)

Voigt RG, Llorente AM, Jensen CL, Fraley JK, Berretta MC, and Heird WC. "A randomized, double-blind, placebo-controlled trial of docosahexaenoic acid supplementation in children with attention-deficit/hyperactivity disorder." J Pediatr 139: 189-196 (2001)

Willatts P, Forsyth JS, DiModugno MK, Varma S, and Colvin M. "Effect of long-chain polyunsaturated fatty acids in infant formula on problem solving at 10 months of age." Lancet 352: 688-691 (19980

Yamada T, Strong JP, Ishii T, Ueno T, Koyama M, Wagayama H, Shimizu A, Sakai T, Malcom GT, and Guzman MA. "Atherosclerosis and omega-3 fatty acids in the populations of a fishing village and a farming village in Japan." Atherosclerosis 153: 469-481 (2000)

Yehuda S, Rabinovitz S, Carasso RL, and Mostofsky DI. "Essential fatty acid preparation improves Alzheimer's patients quality of life." Int J Neurosci 87: 141-149 (1996)

Yehuda S, Rabinovitz S, and Mostofsky DI. "Essential fatty acids are mediators of brain biochemistry and cognitive functions." J Neurosci Res 56: 565-570 (1999)

Zametkin AJ, Nordahl TE, et al. "Cerebral glucose metabolism in adults with hyperactivity of childhood onset." N Engl J Med 323(:1361-1366 (1990)

Zametkin AJ and Ernst M. "Problems in the management of attention-deficit-hyperactivity disorder." N Engl J Med 340: 40-46 (1999)

Zimmer L, Hembert S, Durand G, Breton P, Guillotau D, Besnard JC, and Chalon S. "Chronic n-3 polyunsaturated fatty acid diet-deficiency acts on dopamine metabolism in the rat frontal cortex." Neurosci Letter 240: 177-181 (1998)

Zuijdgeest-van Leeuwen SD, Dagnelie PC, Rietveld T, van den Berg JW, and Wilson JH. "Incorporation and washout of orally adminsistered n-3 fatty acid ethyl esters in different plasma lipid fractions." Br J Nutr 82: 481-488 (1999)